

New York City  
 May 12-15, 1985

Type name, address, and telephone number of author who should receive  
 correspondence in Box F and complete Boxes A, B, C, D and E.

ABSTRACT RECEIPT DEADLINE: JANUARY 11, 1985

TYPE ABSTRACT BELOW. BE SURE TO STAY WITHIN BORDER

(For office use)

NANB WATERBORNE HEPATITIS IN INDIA. B.N.Tandon,  
B.M. Gandhi, Y.K.Joshi, M. Irshad, H.Gupta, Dept.  
 of Gastroenterology, All-India Institute of  
 Medical Sciences, New Delhi-110029, India.

Scarcity and fecal contamination of drinking  
 water are responsible for high prevalence of sever-  
 al viral, bacterial and parasitic diseases in  
 India. We studied the frequency of supposedly wat-  
 erborne hepatitis virus Non-A non-B infection amo-  
 ngst patients of hepatitis at hepatology referral  
 centre in Northern India. Diagnosis of HV-NANB in-  
 fection of HVB due to negative IgM core antibody  
 and HVA due to absence of anti-HAV-IgM in the sera  
 of the patients. Anti HBc-IgM was studied by  
 Cozym-M kit of Abbot Lab, while anti HAV-IgM and  
 HBsAg were investigated by enzyme-linked immuno-  
 sorbent assay. Negative history of blood transfu-  
 sion, fluid infusion and injections and epidemio-  
 logical evidences in favour of fecal contamination  
 of drinking water were used to suggest waterborne  
 instead of post-transfusion HV-NANB infection. The  
 results of study of 778 patients of hepatitis  
 revealed that HV-NANB was the cause of disease in  
 87 % cases of subacute hepatitis, 58 % of acute  
 viral hepatitis, 58 % of fulminant hepatitis and  
 66 % of epidemic hepatitis. 32 % of subacute  
 hepatitis cases had HV-NANB infection in HBsAg  
 carriers, while 9 % of fulminant hepatitis and 6 %  
 each of acute sporadic and epidemic hepatitis e  
 cases were HBsAg carriers with associated HV-NANB  
 infection. Our results suggest that HV-NANB(water-  
 borne) is a major public healthproblem in India. It  
 also leads to more serious and highly fatal forms  
 of subacute and fulminant hepatitis. Associated  
 infection of HV-NANB in HBsAg carriers is danger-  
 ous. Prevention of fecal contamination of drinking  
 water will substantially reduce the incidence of  
 viral hepatitis in India and probably other  
 developing countries.

A

AGA CATEGORIES: Circle one only.

Molecular Biology	Clinical
Morphology/Pathology	Stomach
Hormones	Pancreas
Immunology/Microbiology	Motility
Intestine: absorption, secretion	Oncology

OR

AGA/AASLD CATEGORIES: Circle one only.

Immunology; Reticuloendothelial system  
 Collagen and Fibrosis; Cirrhosis; Transplantation  
 Morphology; Cancer; Regeneration  
 Chronic Hepatitis; Primary Biliary Cirrhosis  
Viral Hepatitis  
 Genetic and Metabolic Liver Disease  
 Complications of Liver Disease; Portal Hypertension  
 Alcohol and the Liver  
 Bile Acids; Biliary Lipids; Gallstones  
 Hepatic Transport; Bile Formation; Cholestasis; Bilirubin  
 Cell Injury and Death; Hepatotoxicity  
 Drug Metabolism and Kinetics

B

**COST OF PUBLICATION**

Payment enclosed (\$35.00)

Check no. \_\_\_\_\_

(payable to AGA)

Payment delayed (\$40.00)

Purchase order no. \_\_\_\_\_

Issued by \_\_\_\_\_

(Institution name)

A copy of this abstract *must* be attached to original  
 purchase order for identification. Payment must be  
 made in U.S. currency.

C

**IMPORTANT**

The principal author affirms that the material  
 herein: 1) will not have been published as an  
 article by May 11, 1985 or have been presented in  
 the U.S. at a national meeting of a national  
 society, or abroad at a major foreign or  
 international meeting; 2) that if human subjects  
 were exposed to risks not required by their medic  
 needs, the study was approved by an appropriate  
 committee or, if no such committee was available  
 and informed consent was needed, it was obtained  
 in accordance with the principles enunciated in  
 "The Institutional Guide to DHEW Policy on  
 Protection of Human Subjects"; or 3) any anima  
 studies conform with the "Guiding Principles in  
 the Care and Use of Animals" of the American  
 Physiological Society.

*B.N. Tandon*

(Signature of Principal Author)

D. If movie projection equipment is needed for oral presentation check here: \_\_\_\_\_

E. If submitted for Student Research Prize competition check here: \_\_\_\_\_

F. Name \_\_\_\_\_

Address \_\_\_\_\_

Telephone, Office: \_\_\_\_\_ Home: \_\_\_\_\_

(area code)

(area code)

ABSTRACT FORM FOR AMERICAN GASTROENTEROLOGICAL ASSOCIATION  
AMERICAN ASSOCIATION FOR THE STUDY OF LIVER DISEASES  
GASTROENTEROLOGY RESEARCH GROUP

New York City  
May 12-15, 1985

Type name, address, and telephone number of author who should receive  
correspondence in Box F and complete Boxes A, B, C, D and E.

ABSTRACT RECEIPT DEADLINE: JANUARY 11, 1985

NANB WATERBORNE HEPATITIS IN INDIA. B.N.Tandon,  
B.M. Gandhi, Y.K. Joshi, M. Irshad, H.Gupta,  
Dept. of Gastroenterology, All-India Institute of  
Medical Sciences, Ansari Nagar, New Delhi, India.

Scarcity and fecal contamination of drinking  
water are responsible for high prevalence of  
several viral, bacterial and parasitic diseases in  
many developing countries of the world, including  
India. We studied the frequency of supposedly  
waterborne hepatitis virus, Non-A non-B infection  
amongst patients of hepatitis at hepatology refer-  
ral centre in Northern India. Diagnosis of HV-NANB  
infection was established by exclusion of acute  
infection of HVB due to negative IgM core antibody  
and HVA due to absence of anti HAV-IgM in the sera  
of the patients. Anti HBc-IgM was studied by  
Corzym-M kit of Abbott Labs, USA while anti HAV-IgM  
and HBsAg were investigated by enzyme-linked  
immunosorbent assay. Negative history of blood  
transfusion, fluid infusion and injections and epi-  
demiological evidences in favour of fecal contami-  
nation of drinking water were used to suggest  
waterborne instead of post-transfusion HV-NANB  
infection. The results of study of 778 patients of  
hepatitis revealed that HV-NANB was the cause of  
disease in 87 % cases of subacute hepatitis, 58 %  
of acute viral hepatitis, 58 % of fulminant hepa-  
titis and 66 % of epidemic hepatitis. 32 % of sub-  
acute hepatitis cases had HV-NANB infection in  
HBsAg carriers, while 9 % of fulminant hepatitis  
and 6 % each of acute sporadic and epidemic hepa-  
titis cases were HBsAg carriers with associated  
HV-NANB infection. Our results suggest that HV-  
NANB(waterborne) is a major public health problem  
in India. It also leads to more serious and highly  
fatal forms of subacute and fulminant hepatitis.  
Associated infection of HV-NANB in HBsAg carriers  
is dangerous. Prevention of fecal contamination of  
drinking water will substantially reduce the  
incidence of viral hepatitis in India and probably  
other developing countries.

A

AGA CATEGORIES: Circle one only

Molecular Biology  
Morphology Pathology  
Hormones  
Immunology Microbiology  
Intestine absorption, secretion

OR

AGA AASLD CATEGORIES: Circle

Immunology, Reticuloendothelial sys  
Collagen and Fibrosis, Cirrhosis, Tran  
Morphology, Cancer, Regeneration  
Chronic Hepatitis, Primary Biliary C  
Viral Hepatitis  
Genetic and Metabolic Liver Disease  
Complications of Liver Disease, Port  
Alcohol and the Liver  
Bile Acids, Biliary Lipids, Gallstones  
Hepatic Transport, Bile Formation, C  
Cell Injury and Death, Hepatotoxicity  
Drug Metabolism and Kinetics

B

COST OF PUBLICATION

Payment enclosed (\$35.00)

Check no. \_\_\_\_\_

(payable to AGA)

Payment delayed (\$40.00)

Purchase order no. \_\_\_\_\_

Issued by \_\_\_\_\_

(Institution)

A copy of this abstract must  
purchase order for identification  
made in U.S. currency.

C

IMPLICATIONS

The principal author affirms  
herein: 1) will not have been  
article by May 11, 1985 in  
the U.S. at a national meeting  
society, or abroad at a national  
international meeting, 2) was  
were exposed to risks not  
needs, the study was approved  
committee or, if no such