

Non-A non-B antigen and antibody detection by agar gel diffusion.

<https://www.ncbi.nlm.nih.gov/pubmed/6818140>

PMID: 6818140

Abstract

Indian J Med Res. 1982 Oct;76:591-3.

Non-A non-B antigen and antibody detection by agar gel diffusion.

Gandhi BM, Joshi YK, Bijlani L, Tandon BN.

Department of Gastroenterology & Human Nutrition, All India Institute of Medical Sciences, Ansari Nagar, New Delhi-110029, India

Abstract

Serum samples from 60 patients of acute viral hepatitis negative for both hepatitis A and B Virus were tested for the presence of non-A non-B hepatitis viral antigens and antibody. Seventeen serum samples from small normal healthy persons were also tested under identical condition. Significant difference was observed for transaminase value in patients positive for antibody and the antigen. Antigen was detected in acute phase sera when SGOT- SGPT levels were significantly elevated.

References

2. Tandon B.N., Joshi, Y.K., Jain, S.K., Gandhi, B.M., Mathisean, L.R., and Tandon, H.D. An epidemic of non A non B hepatitis in north india. Indian J Med. Res. 75 (1982) 739.
7. Tandon, B.N., Krishnamurthy, L. and Gandhi B.M. HBeAg and anti- HBe in acute and chronic liver disease. Indian J Med. Res.70 (1979) 102.