

A dot immunobinding assay (DIA) on nitrocellulose membrane for the serological detection of antibodies to *Entamoeba histolytica*

[https://www.travelmedicinejournal.com/article/0035-9203\(86\)90246-4/pdf](https://www.travelmedicinejournal.com/article/0035-9203(86)90246-4/pdf)

DOI: [https://doi.org/10.1016/0035-9203\(86\)90246-4](https://doi.org/10.1016/0035-9203(86)90246-4)

Abstract

Travel Medicine and Infectious Disease 80(6): 883-885, 1986

A dot immunobinding assay (DIA) on nitrocellulose membrane for the serological detection of antibodies to *Entamoeba histolytica*

B.M. Gandhi

Dept. of Gastroenterology and Human Nutrition, All-India Institute of Medical Sciences, New Delhi 110029, India

Abstract

A rapid, cheap, simple and specific serological test of adequate sensitivity for detecting IgG antibodies against *Entamoeba histolytica* antigen is described. Axenically cultured amoebic antigen was used to precoat the nitrocellulose membrane. The strips were incubated with test samples and later with horseradish peroxidase (HRPO) labelled protein-A conjugate. A dark blue spot was obtained by treatment with peroxidase substrate, 4-chloro-1-naphthol, in positive samples. Serum samples from 32 healthy controls, 45 patients with acute amoebic liver abscess and 10 asymptomatic *E. histolytica* cyst passers were tested. This test was positive in 93% of cases of amoebic liver abscess, 3% of healthy controls and none of the cyst passers; its sensitivity (97%) and specificity (93%) were as good as that of the ELISA test and, because it is simple, quick and cheap it is recommended as a serological test of choice for the diagnosis of invasive amoebiasis.