

Variations in plasma lipid concentration during examination stress

<https://www.ncbi.nlm.nih.gov/pubmed/16250755>

PMID: 16250755, DOI: 10.1207/s15327558ijbm0303_5

Cited in:

Int J Behav Med. 1996;3(3):251-65.

Variations in plasma lipid concentration during examination stress

McCann BS¹, Benjamin GA, Wilkinson CW, Carter J, Retzlaff BM, Russo J, Knopp RH.

¹Cardiovascular Behavioral Medicine Program, Psychiatry and Behavioral Sciences, University of Washington School of Medicine, Seattle 98101-1827, USA. mccann@u.washington.edu

Abstract

The effect of psychological stress on plasma lipids was studied in 40 law students (20 men and 20 women). Plasma cholesterol, triglyceride, and apolipoprotein concentrations were examined at the beginning of the quarter and during the week before final examinations. Cortisol, blood pressure, and heart rate, and self-reports of stress and workload were also measured to verify that examinations were associated with increased stress levels. Perceived stress, perceived workload, and cortisol increased before examinations. Low density lipoprotein cholesterol (LDL-C) increased 5.8 +/- 13.9 mg/dL, and apolipoprotein B (apo B) increased 2.9 +/- 4.0 mg/dL. High density lipoprotein cholesterol decreased in women only. These changes were not due to changes in dietary intake or indexes of plasma volume. However, changes in cortisol and changes in LDL-C and apo B were associated, suggesting a neuroendocrine component to the effects. These results suggest that episodic, stressful situations may lead to potentially atherogenic changes in lipid and lipoprotein concentrations.

Reference

Bijlani RL, Gandhi BM, Tandon BN. Effect of examination stress on serum lipid profile. *Trop Gastroenterol* 1983;4(3):168-70.