

Acute sporadic viral hepatitis in urban population of a tribal district in Madhya Pradesh

medind.nic.in/ibv/t98/i2/ibvt98i2p105.pdf

Cited In:

INDIAN PEDIATRICS , VOLUME 35-FEBRUARY 1998

Acute sporadic viral hepatitis in urban population of a tribal district in Madhya Pradesh

Jagvir Singh, Charu Prakash, R. Panda*, D. Bora, D.C. Jain and K.K. Datta

From the National Institute of Communicable Diseases, 22 Shamnath Marg, Delhi 110 054 and *National Institute of Communicable Diseases, Malaria Research Field Station, Nayapara, Jagdalpur 494 001, Madhya Pradesh.

Reprint requests: Dr. Jagvir Singh, Deputy Director, Epidemiology Division, National Institute of Communicable Diseases, 22 Shamnath Marg, Delhi 110 054.

Manuscript received: July 14,1997; Initial review completed: August 12,1997; Revision accepted: September 24,1997

Abstract

Objective: To estimate the incidence of acute sporadic viral hepatitis and describe its epidemiology in an urban population. Design: A retrospective community survey for jaundice cases. Setting: Headquarter town of a tribal district, Bastar, in Madhya Pradesh state, India. Method: Trained paramedics surveyed about 51,643 population to detect cases of jaundice which occurred in the past one year. Cases were examined to collect clinical and epidemiological data. Blood samples were drawn from all cases who had jaundice in the past 3 months for testing them for markers of viral hepatitis. Results: Study estimated the annual incidence of jaundice cases as 244 (95% CI 201-287) per 100,000 population. Almost 95% jaundice cases occurred in summer and monsoon months. People from all socio-economic strata were affected. The incidence of jaundice was found to be the highest in children below 15 years of age (3.7 per 1000) which declined significantly with the increase in age ($p=0.0000$). The overall incidence in two sexes was not different statistically ($p=0.7$). Of 57 cases who had jaundice in the past 3 months, 19 (33%) were confirmed as having viral hepatitis. Hepatitis A and E combined together contributed 68% (13/19) of acute sporadic cases of viral hepatitis, whereas hepatitis B, C and D accounted for the remaining 32% of the cases. Conclusion: The study found the annual incidence of laboratory supported cases of viral hepatitis to be 81 (95% CI 57-106) per 100,000 population, which suggests that it is an important public health problem in India. Hepatitis A was much more prevalent than hepatitis E. Etiology of almost two-thirds of jaundice cases could not be established which require further community studies

Reference

2. Tandon BN, Gandhi BM, Joshi YK. Etiological spectrum of viral hepatitis and prevalence of markers of hepatitis A and B infection in north India. Bull World Health Organ 1984; 62: 67-73.

3. Tandon BN, Gandhi BM, Joshi YK, Irshad M, Gupta H. Hepatitis virus non-A, nonB: The cause of major public health problem in India. Bull World Health Organ 1985; 63: 931-934.