

Hepatitis e virus infection in fulminant hepatitis patients and an apparently healthy population in bangladesh

BANGLADESH www.ajtmh.org/content/66/6/721.full.pdf

Cited In :

Am. J. Trop. Med. Hyg., 66(6), 2002, pp. 721–724

Copyright © 2002 by The American Society of Tropical Medicine and Hygiene

Hepatitis e virus infection in fulminant hepatitis patients and an apparently healthy population in bangladesh

Aleemuzzaman Sheikh, Masahiko Sugitani, Noriko Kinukawa, Mitsuhiko Moriyama, Yasuyuki Arakawa, Kazuo Komiyama, Tian-Cheng Li, Naokazu Takeda, Shamsuddin M. Ishaque, Mahmud Hasan, And Koyu Suzuki

Department of Pathology and Third Department of Internal Medicine, Nihon University School of Medicine, Tokyo, Japan; Department of Oral Pathology, Nihon University School of Dentistry, Tokyo, Japan; Department of Virology II, National Infectious Institute, Tokyo, Japan; Department of Diagnostic Pathology, Saint Luke's International Hospital, Tokyo, Japan; Department of Gastroenterology, Bangabandhu Sheikh Mujib Medical University, Dhaka, Bangladesh

Abstract

This is the first study comparing hepatitis E virus (HEV) infection in Bangladesh in fulminant hepatitis (FH) patients presumed to have a viral cause and in the apparently healthy population. Sera from 22 FH patients were analyzed for antibodies to hepatitis A virus (HAV), hepatitis B virus (HBV), hepatitis C and D viruses, and HEV and for hepatitis B surface antigen (HBsAg). Anti-HEV immunoglobulin M (IgM) was detected in the sera of 63.6% of patients, whereas 35.7% were positive for HBsAg. A high prevalence of HEV infection (83.3%) was noted in the HBV carriers. Serum samples from 273 apparently healthy individuals were tested for antibodies to HAV and HEV. AntiHEV IgM was detected in 7.3% of the samples. The seroprevalence of HAV differed from that of HEV in the same population because all samples were negative for anti-HAV IgM. These data indicate that HEV infection is highly endemic in Bangladesh.

Reference

23. Tandon BN, Gandhi BM, Joshi YK, 1984. Etiological spectrum of viral hepatitis and prevalence of markers of hepatitis A and B virus infection in north India. Bull WHO 62: 67–73.
34. Gandhi BM, Joshi YK, Tandon BN, 1981. Prevalence of virus A exposure and viral hepatitis A in India. Lancet ii: 374.