

Serological responses to hepatitis B virus infection in multi-transfused thalassemic children

<https://europepmc.org/abstract/med/1601501>

Cited in:

Indian Pediatrics 29(1):73-77, 1992

Serological responses to hepatitis B virus infection in multi-transfused thalassemic children.

Gulati S, Marwaha RK, Dilawari JB, Midha U, Walia BN

Department of Pediatrics, Postgraduate Institute of Medical Education and Research, Chandigarh.

Abstract

One hundred children with beta-thalassemia major were studied prospectively. A onetime analysis of serum samples was carried out for a battery of hepatitis B viral markers viz., HBsAg, anti-HBs and anti-HBc. Seven mutually different serological patterns were observed. The commonest profile seen in 49 patients was a combined seropositivity for anti-HBc and anti-HBs indicating past HBV infection with persisting immunity. Definite evidence of active HBV infection (seropositivity for HBsAg and/or HBeAg) was demonstrated in 10 cases, six of these were HBsAg positive. Anti-HBc positivity alone was detected in 17 patients. The remaining 24 children were seropositive for anti-HBs alone suggesting a possible passive transmission of anti-HBs through blood transfusion.

Reference

Patwari SI, Irshad M, Gandhi BN, Joshi YK, Nundy S, Tandon BN. Post-transfusion hepatitis — a prospective study. *Indian J Med Res* 1986; 84:508–10.