

Significance of anti-pre-S antibodies in patients with fulminant hepatic failure

<https://link.springer.com/article/10.1007/BF02779341>

<https://doi.org/10.1007/BF02779341>

Cited In:

Gastroenterologia Japonica 25(4): 499-502, 1990

Significance of anti-pre-S antibodies in patients with fulminant hepatic failure

M. Irshad¹, B. M. Gandhi¹ S. K. Acharya¹ Y. K. Joshi¹ B. N. Tandon¹

¹Department of Gastroenterology and Human Nutrition All India Institute of Medical Sciences, New Delhi India

Summary

Anti-pre-S antibody was tested in 38 sera from patients with fulminant hepatitis (positive for HBsAg and/or IgM anti-HBc) using a specific solid phase enzyme linked immunosorbent assay (ELISA). Anti-pre-S activity was detected in 50 percent sera samples positive for HBsAg but negative for IgM antiHBc. There were 12.5% sera positive for both HBsAg as well as IgM anti-HBc and 75% sera negative for HBsAg but positive for IgM anti-HBc. The prevalence of HBV-specific DNA-polymerase activity was high in all the three groups whereas anti-HBs positivity was low. Anti-pre-S activity was observed both in the presence as well as in the absence of DNA-polymerase activity. High-anti-pre-S level in fulminant hepatitis B patients was assumed to be implicated in the fast clearance of HBsAg from circulation.

Key Words: anti-pre-S ELISA fulminant hepatitis HBsAg clearance

Reference

Irshad M, Gandhi BM, Acharya SK, et al: Anti-pre-S antibodies in different groups of patients with hepatitis B virus (HBV) infection. *J Gastroenterol Hepatol* 1989;4:25–32

Irshad M. Gandhi BM. Chawla TC, et al: Studies on HBsAg binding with polymerised human serum albumin by ELISA. *J Virol Meth* 1987;16:75–85