

## **Prognostic value of HBsAg/IgM complexes in hepatitis B patients: nature of the proteins involved.**

<https://www.ncbi.nlm.nih.gov/pubmed/3171555>

### **Cited in:**

J Med Virol. 1988 Jul;25(3):309-15.

## **Prognostic value of HBsAg/IgM complexes in hepatitis B patients: nature of the proteins involved.**

Grangeot-Keros L<sup>1</sup>, Pelletier G, Briantais MJ, Pillot J.

<sup>1</sup> Inserm U 131 Hôpital Antoine Bécclère, Clamart, France.

### **Abstract**

Hepatitis B surface antigen (HBsAg)/IgM complexes were measured using a solid phase radioimmunoassay in sera of patients with acute or chronic hepatitis B. These complexes were found in 6 out of 18 patients four weeks after the onset of the disease and only one of them developed chronic hepatitis. HBsAg/IgM complexes correlated with HBsAg, hepatitis B e antigen (HBeAg), and pre-S2 concentration. The precipitation of HBsAg/IgM reactivity by polyethylene-glycol (PEG) and the binding of this activity to the surface of certain uncoated enzyme linked immunosorbent assay (ELISA) plates indicates that HBsAg/IgM positivity may reflect the presence of circulating complexes in serum. HBsAg and pre-S2 were found as components of the complexes but anti-HBs, anti-pre-S2, and polymerized human serum albumin (pHSA) were not. An immune binding between HBsAg and IgM is still questionable. Whatever the nature of the HBsAg/IgM complexes, their detection does not seem to be an earlier indicator of prognosis than HBeAg and/or pre-S2.

### **Reference**

M. Irshad, B.M. Gandhi, T.C. Chawla, S.K. Acharya, Y.K. Joshi, B.N. Tandon, Studies on HBsAg binding with polymerised human serum albumin by ELISA, J. Virol. Methods 16 (1987) 75-85.